


IDENTIFYING RISKS AND IMPROVING OUTCOMES FOR ADOLESCENT PATIENTS

High-Risk Behaviors

High-risk behaviors are the **primary causes of morbidity and mortality** in adolescent patients (ages 12 to 21):¹

- » **Substance abuse**
- » **Unsafe sexual activity**
- » **Interpersonal violence**
- » **Suicide**



Why Confidentiality Matters

- Adolescents are **more likely to discuss high-risk behaviors** if they believe their care is confidential.^{2,4,5}
- Adolescents **answer confidential screenings more honestly**.⁶
- State and national **laws allow minors to receive confidential care** related to sexual health, mental health, and substance abuse.

Barriers to Confidential Care



- There is **low knowledge about minor consent laws**.^{7,8,9}
- Less than half of adolescents receive a yearly well or preventative exam. **Most do not spend any time alone with their provider** during that visit.¹⁰
- Providers have noted a **lack of expertise, insurance issues, and concerns about medical records**.¹¹

Advantages of Screening Tools

- Screening tools **provide a comprehensive picture** of the patient.
- They **increase efficiency and effectiveness of care**, allowing physicians to tailor their conversations with patients.
- When paired with effective counseling and intervention, **they can make a significant impact on adolescent high-risk behaviors**.¹²

M | **ADOLESCENT HEALTH INITIATIVE**
UNIVERSITY OF MICHIGAN HEALTH SYSTEM

Example of a Confidential Work Flow

- 1** At check-in, front desk staff gives parent/guardian and patient a letter about confidential time with adolescent patients.
- 2** Medical Assistant (MA) calls patient, explains to parent/guardian, "I'll be bringing your child back to get their vital signs and have them complete a brief health survey. Then I'll bring you to the room."
- 3** MA places patient in an exam room, has them complete the screening tool, brings the results to the provider to review, and then brings back parent/guardian.
- 4** Provider meets with parent/guardian and patient, and then asks the parent/guardian to step out for confidential time. Provider then discusses the risk screen confidentially with the patient.

REFERENCES: **1)** US Congress, Office of Technology Assessment. Adolescent Health: Summary and Policy Options. Washington DC: US Government Printing Office, 1991. OTA-H-468; **2)** Bethell C, Klein J, Peck C. Assessing health system provision of adolescent preventive services: the Young Adult Healthcare Survey. Med Care. 2001. 39(5):478-490; **3)** Blum RW, Beuhring T, Wunderlich M, Resnick MD. Don't ask, they won't tell: the quality of adolescent health screening in five practice settings. Am J Public Health. 1996. 86:1767-1772; **4)** Klein JD, Wilson KM. Delivering Quality Care: Adolescents' discussion of health risks with their providers. J Adolesc Health. 2002. 30:190-195; **5)** Ford CA, Bearman PS, Moody J. Foregone health care among adolescents. JAMA. 1999. 282(23):2227-2234; **6)** Kadivar, H., Thompson, L., Wegman, M. Adolescent views on comprehensive health risk assessment and counseling: Assessing gender differences. Journal of Adolescent Health. 2014. 55: 24-32.; **7)** Rock EM, Simmons PS. Physician knowledge and attitudes of Minnesota laws concerning adolescent health care. J Pediatr Adolesc Gynecol. 2003. 16:101-108; **8)** Loertscher L, Simmons PS. Adolescents' knowledge of and attitudes toward Minnesota laws concerning adolescent medical care. J of Pediatr Adolesc Gynecol. 2006. 19:205-207; **9)** Cutler EM, Bateman MD, Wollan PC, Simmons PS. Parental knowledge and attitudes of Minnesota laws concerning adolescent medical care. Pediatrics. 1999. 103:582-587; **10)** Irwin CE Jr, Adams SH, Park MJ, Newacheck PW. Preventive care for adolescents: few get visits and fewer get services. Pediatrics. 2009. 123(4):e565-572; **11)** Riley M, Ahmed S, Reed BD, Quint EH. Physician Knowledge and Attitudes around Confidential Care for Minor Patients. J Pediatr Adolesc Gynecol. 2015. 28(4): 243-9; **12)** Ozer EM, Adams, SH, Orrell-Valente JK, et al. Does Delivering Preventative Services in Primary Care Reduce Adolescent Risky Behavior? Journal of Adolescent Health. 49(5):476-482.